



TITLE:

Effects of trunk lean and foot lift exercises in sitting position on abdominal muscle activity and the contribution rate of transversus abdominis

AUTHOR(S):

Motomura, Yoshiki; Tateuchi, Hiroshige;
Komamura, Tomohito; Yagi, Yuta; Nakao, Sayaka;
Ichihashi, Noriaki

CITATION:

Motomura, Yoshiki ...[et al]. Effects of trunk lean and foot lift exercises in sitting position on abdominal muscle activity and the contribution rate of transversus abdominis. *European Journal of Applied Physiology* 2021, 121(1): 173-181

ISSUE DATE:

2021-01

URL:

<http://hdl.handle.net/2433/261777>

RIGHT:

This is a post-peer-review, pre-copyedit version of an article published in *European Journal of Applied Physiology*. The final authenticated version is available online at: <http://dx.doi.org/10.1007/s00421-020-04508-0>; The full-text file will be made open to the public on 30 September 2021 in accordance with publisher's 'Terms and Conditions for Self-Archiving'; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。; This is not the published version. Please cite only the published version.

1 **Effects of trunk lean and foot lift exercises in sitting position on abdominal muscle activity**
2 **and the contribution rate of transversus abdominis**

3

4 **Yoshiki Motomura¹, Hiroshige Tateuchi¹, Tomohito Komamura², Yuta Yagi³, Sayaka Nakao¹, Noriaki**
5 **Ichihashi¹**

6

7 1. Human Health Sciences, Graduate School of Medicine, Kyoto University, Kyoto, Kyoto, Japan

8 2. Division of Rehabilitation Medicine, Chiba University Hospital, Chiba, Chiba, Japan

9 3. Department of Rehabilitation, Rinku General Medical Center, Izumisano, Osaka, Japan

10

11 **Corresponding author**

12 Yoshiki Motomura (E-mail: motomura.yoshiki.32z@kyoto-u.jp)

13

14 **ORCID**

15 Yoshiki Motomura: 0000-0002-6544-0678

16 Sayaka Nakao: 0000-0001-5714-0336

17 Noriaki Ichihashi: 0000-0003-2508-2172

18

19

20 **Acknowledgements**

21 The authors gratefully acknowledge all participants involved in this study. This study was not funded by any
22 institutions, agencies, or companies.

23

24 **Declarations**

25 **Funding:** Not applicable.

26 **Conflicts of interest:** Not applicable.

27 **Ethics approval:** All the procedures performed in the studies involving human participants were in accordance
28 with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki
29 declaration and its later amendments or comparable ethical standards. This study was approved by the ethics
30 committee of Kyoto University Graduate School and the Faculty of Medicine (R0546-2)

31 **Consent:** Informed consent was obtained from all individual participants involved in the study.

32 **Data and/or Code availability:** All data generated or analysed during this study are included in this published
33 article.

34 **Authors' contribution statements:** YM, HT, SN, and NI conceived and designed the research. YM, TK, and YY
35 conducted the experiments. YM, HT, and NI analyzed the data. YM, HT, SN, and NI wrote the manuscript. All
36 the authors have read and approved the manuscript.

37

38

39 **Abstract**

40 *Purpose:* Abdominal hollowing exercise has been recommended to improve trunk stability. Trunk lean and foot
41 lift exercises while sitting may easily promote abdominal muscle activity even in people who cannot perform
42 abdominal hollowing consciously. The purpose of the present study was to examine the changes in abdominal
43 muscle activity and contribution rate of the transversus abdominis muscle (TrA) when leaning the trunk and lifting
44 the foot during sitting.

45 *Methods:* The muscle stiffnesses (indicators of muscle activity) of the right rectus abdominis, external oblique,
46 internal oblique, and TrA of 14 healthy men were measured during abdominal hollowing and the following nine
47 sitting tasks: reference posture, 15° and maximal posterior trunk lean, 20° and maximal ipsilateral and contralateral
48 trunk lean, and ipsilateral and contralateral foot lift. The TrA contribution rate was calculated by dividing the TrA
49 stiffness by the sum of the abdominal muscles' stiffnesses.

50 *Results:* The TrA stiffness was significantly higher in abdominal hollowing than in reference posture, posterior and
51 ipsilateral trunk lean, and ipsilateral foot lift, but not higher than in contralateral trunk lean and contralateral foot
52 lift. There was no significant difference in the TrA contribution rates between abdominal hollowing and ipsilateral
53 or contralateral foot lift.

54 *Conclusion:* The contralateral trunk lean or contralateral foot lift could enhance TrA activity for people who cannot
55 perform abdominal hollowing consciously. The contralateral foot lift could particularly be beneficial to obtain
56 selective activity of TrA.

57

58 **Keywords**

59 abdominal hollowing, muscle stiffness, transversus abdominis, internal oblique, external oblique, rectus abdominis

60

61 **Abbreviations**

62 TrA Transversus abdominis muscle

63 ANOVA Analysis of variance

64 SWE Shear wave elastography

65

66

67 Introduction

68 The transversus abdominis muscle (TrA) plays an important role in trunk stabilization while moving the
69 extremities (Hodges and Richardson 1996, 1998; Hodges et al. 1997; Okubo et al. 2013). Since the TrA acts to
70 tighten the abdomen even when the activities of the other abdominal muscles remain unchanged, greater TrA
71 activity may allow for a more effective increase in intra-abdominal pressure, which increases the stiffness of the
72 lumbar spine (Hodges et al. 2005). Therefore, improving TrA contribution rate, which is the percentage of TrA
73 activity in all the abdominal muscle activities, is required to increase spinal stiffness and reduce spinal loading
74 (Aspden 1988).

75 Abdominal hollowing exercise, which retracts the abdomen consciously, has been commonly used to train the
76 TrA (Beith et al. 2001; Koh et al. 2014). Isolated TrA activation using very low-intensity abdominal hollowing
77 may be effective to promote muscle recruitment such as improving the delay in neuromuscular activity of TrA
78 (Tsao and Hodges 2007). On the other hand, a previous study found that as the intensity of abdominal hollowing
79 increased, the TrA activity increased significantly and the ratio of the TrA to the internal oblique, external oblique,
80 and rectus abdominis did not change (Shimizu et al. 2019). That is, abdominal hollowing at a higher intensity may
81 more effectively improve the function of the TrA that stabilizes the trunk.

82 Greater decrease in the abdominal cavity during abdominal hollowing reflects stronger contraction of the TrA
83 (Richardson et al. 2004). Hides et al. (2008) reported that there was no significant difference in the TrA thickness
84 and abdominal cavity at rest between those with and without low back pain, and the abdominal cavity during
85 abdominal hollowing was significantly larger in those with low back pain than those without. Therefore, patients

86 with low back pain may have difficulty exerting voluntary TrA contraction even in the absence of atrophy. Hence,
87 training methods targeting involuntary activation of TrA are important for patients with low back pain.

88 The prone bridge exercise activates abdominal muscles involuntarily by resisting the gravity from the posture
89 change (Okubo et al. 2010; Shiju Majeed et al. 2019). However, methods promoting abdominal muscle activity
90 through dynamic posture changes, such as prone bridge, have high physical loads and are not necessarily safe for
91 patients with low back pain (Ekstrom et al. 2008; Bhadauria and Gurudut 2017). Though some studies have
92 reported the relation between abdominal muscle activity and sagittal spinal alignment in sitting (O’Sullivan et al.
93 2002; Astfalck et al. 2010; Claus et al. 2018), these studies did not focus on exercises. However, considering these
94 studies, the TrA activity may be involuntarily increased by leaning the trunk or lifting the foot during sitting, even
95 in patients with low back pain and elderly people with difficulty in changing posture dynamically with high
96 intensity. Foot lift exercises are not changed trunk posture, but may increase abdominal muscle activity to increase
97 lumbar and pelvic stiffness, in order to stabilize the pelvis and to exert hip flexion torque effectively. Revealing
98 how the abdominal muscles activate when leaning the trunk and lifting the foot during sitting may provide
99 knowledge for rehabilitation to stabilize trunks in patients with low back pain and elderly people.

100 The purpose of this study was to verify the effect of trunk lean and foot lift exercises during sitting on abdominal
101 muscle activity and TrA contribution rate. The hypothesis was that the activity of all abdominal muscles will be
102 highest in the posterior trunk lean because the spine is more unstable in flexion and extension than in lateral flexion
103 (Yamamoto et al. 1989). It was also hypothesized that TrA contribution rate would be highest in the contralateral
104 trunk lean where rectus abdominis activity may be more decreased among the abdominal muscles, according to

105 previous studies (Masani et al. 2009; Eriksson Crommert et al. 2017).

106

107 **Methods**

108 **Participants**

109 A total of 14 healthy men (age, 24.6 ± 2.9 years; height, 172.5 ± 6.1 cm; mass, 66.9 ± 9.0 kg) volunteered for this
110 study. The exclusion criteria were a history of low back pain lasting more than three months(Chou et al. 2007),
111 operation and neurological or orthopedic diseases in the trunk or lower limbs. A power analysis with an α error =
112 0.05, power = 0.80, and effect size $f = 0.25$ (medium) was performed by the G*Power 3.1 analysis software
113 (Heinrich Heine University, Duesseldorf, Germany) for one-way repeated measures analysis of variance (ANOVA).
114 This produced a minimum total sample size of 12. This study was approved by the ethics committee of Kyoto
115 University Graduate School and the Faculty of Medicine (R0546-2) and was conducted in compliance with the
116 Declaration of Helsinki. All participants were provided written informed consent after being briefed with the
117 objectives and the risks involved in the experiment.

118

119 **Experimental protocol**

120 To minimize the differences in muscle activity due to different spinal alignments in each participant's natural sitting
121 position, a reference posture was defined (Fig 1. a). This is the upright sitting posture, whereby the axis from ear
122 lobe to the floor lies between the anterior and posterior superior iliac spine on the sagittal plane. Further visual
123 verification was done by two of our physiotherapists to ensure no remarkable spinal curvature (e.g. thoracic or

124 lumbar hyperflexion). Participants randomly performed tasks maintaining the following postures (Fig 1. b-f):
125 leaning the trunk posterior to 15° and maximum from reference posture (posterior trunk lean), leaning the trunk at
126 20° and maximum to ipsilateral and contralateral from reference posture (ipsilateral and contralateral trunk lean),
127 and lifting the ipsilateral and contralateral foot about 1 cm from the floor (ipsilateral and contralateral foot lift).
128 Participants received feedback from a mirror placed 1.5-m in front of them, and were instructed to perform tasks
129 without trunk flexion/extension, lateral flexion, or rotation. The measurements were conducted while one examiner
130 confirmed there was no obvious deviation of posture during the tasks. Then the participants performed abdominal
131 hollowing with maximal effort in supine position without moving the trunk and pelvis (Fig 1. g). Lumbar lordosis
132 during abdominal hollowing was confirmed by participants using the Stabilizer Pressure Biofeedback unit (PBU,
133 Chattanooga Group, Australia) placed under the lumbar spine, with a constant pressure of 40 mmHg. This was
134 done to standardize pelvic inclination among participants during the maneuver. They were instructed to perform
135 abdominal hollowing while trying to maintain the pressure at 40 mmHg.

136

137 **Shear wave elastography**

138 In each task, muscle stiffnesses of the right TrA, internal oblique, external oblique, and rectus abdominis were
139 measured three times. The measurement sites were determined based on previous studies (Shimizu et al. 2019):
140 TrA and internal oblique muscles, 2-cm medial the anterior superior iliac spine; external oblique, 2.5-cm medial
141 from the point on the axillary line at navel height; and rectus abdominis, 4-cm lateral the navel (Fig 2). Muscle
142 stiffness was calculated using the following formula by shear wave elastography (SWE) mode (musculoskeletal

143 preset) of the Aixplorer ultrasound scanner (v6.4; Supersonic Imagine, Aix-en-Provence, France):

144
$$\mu \text{ (kPa)} = \rho Vs^2,$$

145 where ρ = muscle tissue density (1,000 kg/m³), and Vs = propagation velocity of the shear wave generated by the
146 ultrasonic transducer. An ultrasonic probe (SL15-4 transducer) was in parallel to the fiber orientation of the target
147 muscle. Muscle stiffness was calculated in a 3-mm diameter Q-box at the center of the region of interest placed at
148 the center of each muscle (Fig 2). Reports state that muscle stiffness increases with muscle activity (Bouillard et
149 al. 2011), and there is high reliability of abdominal muscle stiffness measured using SWE (MacDonald et al. 2016;
150 Shimizu et al. 2019). Muscle stiffness was calculated as an average of three measurements for each muscle. After
151 calculating intra-rater reliability (ICC_{1,3}) of these three measurements per task, the reliability of each muscle
152 stiffness was “almost perfect”: TrA, 0.93–1.00; internal oblique, 0.98–1.00; external oblique, 0.98–0.99; and rectus
153 abdominis, 0.93–1.00. The TrA contribution rate was calculated by dividing TrA stiffness by the sum of the
154 stiffnesses of all four abdominal muscles.

155

156 **Spinal and pelvic alignment**

157 Another examiner who did not operate the ultrasonic equipment carefully checked visually to ensure no obvious
158 trunk motion during the task. To verify the degree of spinal flexion and extension, sagittal spinal alignment was
159 measured twice using the Spinal Mouse (Index Ltd., Tokyo, Japan) before every measurement for muscle stiffness.
160 The intra-rater reliabilities (ICC_{1,1}) were then calculated. In 12 participants, excluding 2 with data loss, ICC_{1,1} of
161 spinal alignment data (i.e., the sum of segmental angles from Th1/2 to L5/S) (Tateuchi et al. 2018) ranged from

162 0.73 to 0.88. The average angles of thoracic kyphosis and lumbar lordosis were calculated from these data. The
163 average angle of pelvic posterior inclination at the height of the second sacrum measured three times using an
164 inclinometer (Wixey, USA) was calculated, and intra-rater reliability ($ICC_{1,1}$) ranged from 0.89 to 0.98. The
165 average angle of the maximum spine inclination to posterior and right/left measured three times using a goniometer
166 was calculated.

167

168 **Statistical analysis**

169 Statistical analysis was performed using SPSS version 22.0 (SPSS Japan Inc., Tokyo, Japan). The one-way
170 repeated-measures ANOVA analysis was used to compare the paired datasets between tasks and to investigate
171 whether specific abdominal muscle stiffness or TrA contribution rates would differ depending on the task. When
172 a significant difference was observed, multiple comparisons corrected by the Holm method were performed as a
173 post-hoc test. Dunnet's test was performed to compare the thoracic kyphosis, lumbar lordosis, and pelvic
174 inclination angles between reference posture and other sitting tasks. Additionally, in order to examine the variation
175 among participants, the Pearson correlation analysis was conducted to determine the relationship between TrA
176 contribution rates in each task and the stiffness of the internal oblique, external oblique, and rectus abdominis in
177 the reference posture. A P value <0.05 was considered statistically significant.

178

179 **Results**

180 The muscle stiffness for each muscle in the various tasks is shown in Table 1. All muscle stiffnesses showed

181 significant main effects of tasks in one-way repeated measures ANOVA. TrA stiffness was significantly higher in
182 abdominal hollowing than in all other tasks, except for contralateral trunk lean (at 20° and maximum) and foot lift.
183 TrA stiffness in the maximum contralateral trunk lean was significantly higher than that in the reference posture,
184 posterior trunk lean (at 15° and maximum), and ipsilateral foot lift. The stiffness of the internal oblique was
185 significantly higher in abdominal hollowing than in all other tasks, except for contralateral trunk lean (at 20° and
186 maximum), and was significantly higher in the maximum contralateral trunk lean than in reference posture,
187 posterior trunk lean (at 15° and maximum), ipsilateral trunk lean (at 20° and maximum), and ipsilateral foot lift.
188 The stiffness of the external oblique was significantly higher in the posterior trunk (at 15° and maximum) and
189 contralateral trunk leans (at 20° and maximum) than in all other tasks, but there were no significant differences
190 among the four tasks of the posterior trunk (at 15° and maximum) and contralateral trunk leans (at 20° and
191 maximum). The stiffness of rectus abdominis was significantly higher in the posterior trunk lean at maximum than
192 in all other tasks.

193 The TrA contribution rates in the various tasks is shown in Table 1. There was a significant main effect of task
194 in one-way repeated measures ANOVA. The TrA contribution rate in abdominal hollowing was significantly higher
195 than that in the posterior trunk lean (at 15° and maximum), ipsilateral trunk lean at maximum, and contralateral
196 trunk lean (at 20° and maximum). There was no significant difference in TrA contribution rate between abdominal
197 hollowing and reference posture, ipsilateral trunk lean at 20°, and ipsilateral and contralateral foot lift.

198 The results of thoracic kyphosis angle, lumbar lordosis angle, pelvic inclination angle, and maximum spinal
199 inclination angle are shown in Table 2. The thoracic kyphosis and lumbar lordosis angles were not significantly

200 different between reference posture and other sitting tasks. The pelvic posterior inclination angle was significantly
201 higher in the posterior trunk lean than in reference posture.

202 The additional Pearson correlation analysis showed that the TrA contribution rate in those with high external
203 oblique stiffness in the reference posture tended to be low in the ipsilateral foot lift ($r = -0.742$, $p = 0.002$) and high
204 during maximum abdominal hollowing ($r = 0.519$, $p = 0.057$).

205

206 **Discussion**

207 The present study was the first, to our knowledge, to investigate noninvasively the effects of trunk lean and foot
208 lift exercises during sitting on abdominal muscle activity. High TrA activity was exerted in the contralateral trunk
209 lean and contralateral foot lift during sitting, and the TrA contribution rate in the contralateral foot lift was a similar
210 level to that in maximum abdominal hollowing. These exercises can be performed in elderly people and patients
211 with low back pain, who have difficulty with consciously contracting abdominal muscles such as abdominal
212 hollowing. Our results have elucidated the specific exercises which maximize the activation of TrA and improve
213 TrA contribution rate. Therefore, these may be useful in the consideration of targeted TrA exercises to stabilize the
214 trunk of elderly people and patients with low back pain.

215 Although the TrA activity was highest in abdominal hollowing, TrA activity in the contralateral trunk lean during
216 sitting showed no significant difference to that in abdominal hollowing and tended to be higher than that in
217 reference posture, posterior trunk lean, and ipsilateral foot lift. These results differed from our hypothesis that
218 higher TrA activity will be exerted in the posterior trunk lean because the spine is more unstable in flexion and

219 extension than in lateral flexion (Yamamoto et al. 1989). The TrA may have an important role holding the trunk
220 and maintaining the posture predictively while other muscles contract (Hodges and Richardson 1997; Allison et
221 al. 2008). On the other hand, previous study showed using wire electromyography that the activity of the TrA and
222 internal oblique increased when pulled to contralateral sides, while the activity of the external oblique and rectus
223 abdominis increased when pulled posteriorly (Eriksson Crommert et al. 2017). This study supports our results.
224 Therefore, the present study indicates that all abdominal muscles, even the TrA working to stabilize the trunk, may
225 be specifically activated in postures with external moments in the opposite direction to their anatomical
226 orientations. Moreover, the neutral zone, which is the range of inter-vertebral motion whereby spinal stiffness (i.e.
227 the force required to make a constant displacement between the vertebrae) is the lowest (Panjabi 1992), has been
228 reported to increase with ligament damage and disc degeneration (Panjabi et al. 1989; Hasegawa et al. 2008).
229 Busscher et al. (2009) indicated that the lumbar vertebrae had less spinal stiffness in lateral bending in a wider
230 range of motion than the lower thoracic vertebrae and might have less resistance of passive tissue such as ligaments.
231 Therefore, TrA activity is more likely to increase in lateral trunk lean than posterior trunk lean due to its anatomical
232 function. The present study supports the role of TrA in increasing spinal stiffness. However, because this study did
233 not verify the load on the spine during the task, further studies should determine whether direction-specific activity
234 of the TrA reflects direction-specific properties of the spine.

235 The TrA contribution rate was significantly higher in the foot lift than in the posterior or the contralateral trunk
236 lean, which differed from our hypothesis. This may be because the stiffness of the lumbar spine and pelvis
237 increased with TrA activity (Tesh et al. 1987), making it easier to exert muscle strength of the hip flexors during

238 foot lift. The reason why the activity of the rectus abdominis and oblique abdominal muscles, which are the global
239 muscles (Bergmark 1989), did not increase much may be because the trunk load from gravity was lower in foot
240 lift than in contralateral trunk lean. Therefore, the increase in TrA contribution rate in foot lift may be attributed to
241 these circumstances. On the other hand, the low TrA contribution rate during contralateral trunk lean may be due
242 to the requirement to stabilize not only the lumbopelvic region but also the entire spinal alignment against gravity,
243 rendering isolated TrA activity insufficient. In other words, the rectus abdominis, external oblique and internal
244 oblique muscles may have been activated to stabilize the thorax.

245 TrA acts to tighten the abdomen. It is, however, a thin muscle, therefore is independently not adequate to
246 contribute to spinal stiffness. It is hence suggestive that TrA plays a supportive role in helping the activities of
247 other abdominal muscles. Therefore, high TrA contribution rate (i.e. higher TrA activity when those of other
248 abdominal muscles remain unchanged) may be important in allowing for more effective increase of intra-
249 abdominal pressure, which leads to the increase of spinal stiffness (Hodges et al. 2005; Hides et al. 2006). However,
250 a recent Cochrane review about nonspecific low back pain reported that there were no differences in the effect on
251 improving disability due to low back pain between the specific training for TrA and multifidus muscles and general
252 trunk exercises such as stretching and resistance training (Saragiotto et al. 2016). This is believed to be due to
253 diversity of potential causes of nonspecific back pain (Kiesel et al. 2007). Thus, specific training of the TrA may
254 not necessarily be important for all low back pain patients. In the present study, the variation in the degrees of
255 abdominal muscle stiffness among participants may have affected our results. The additional Pearson correlation
256 analysis have verified the relationship between the TrA contribution rates in each task and the stiffness of the

257 internal oblique, external oblique and rectus abdominis muscles in the reference posture. The results showed that
258 TrA contribution rate in those with high stiffness of external obliques in the reference posture tended to be low
259 during ipsilateral foot lifting ($r = -0.742$) and high during maximum abdominal hollowing ($r = 0.519$). This suggest
260 that the particular exercises required to improve TrA contribution rate may differ according to the properties of
261 abdominal muscles during the sitting position. Further study should better understand which subgroups of patients
262 with low back pain require exercise with a high TrA contribution rate (Hill et al. 2008; Macedo et al. 2014).

263 In this study, characteristics of abdominal muscles were investigated using SWE. Since measurement values of
264 muscle stiffness in this study were similar to those in a previous SWE study (Shimizu et al. 2019), verification of
265 abdominal muscle activities using abdominal muscles' stiffnesses is considered appropriate. Neuromuscular
266 activity measured by a surface or wire electromyography and muscle thickness by an ultrasonic device have been
267 commonly used to verify abdominal muscle activity. However, abdominal muscle thickness changes during
268 contraction may not necessarily be proportional to increases in abdominal muscle activities (Hodges et al. 2003;
269 Whittaker et al. 2013). In addition, surface electromyography cannot measure the TrA, a deep muscle, and wire
270 electromyography is invasive. The SWE in the present study can measure a deep muscle noninvasively and may
271 be useful for verifying abdominal muscle (especially TrA) activity.

272 This study had some limitations. First, spinal lateral flexion and rotation could not be evaluated objectively.
273 Since spinal motion greatly influences abdominal muscle activity because of abdominal muscle anatomy, the
274 experiment paid attention to spinal motion. To avoid fatigue due to an increase in the number of tasks measured,
275 only spinal mobilities in flexion and extension were measured by the Spinal Mouse. However, there were no

276 significant differences in thoracic kyphosis, lumbar lordosis, and pelvic inclination angles between tasks; thus,
277 evident spinal motion probably did not occur in this study. The second limitation was that only men participated
278 in the present study. The mobilities of and load on the sacroiliac joint are reported to be greater in women than in
279 men (Joukar et al. 2018); therefore, since lower fibers of the TrA increase the stiffness of the sacroiliac joint, results
280 may differ in a female study population. Third, the tasks used in present study were not exercises whereby TrA
281 was activated in isolation. Lastly, they may not be appropriate for all patients with low back pain.

282

283 **Conclusion**

284 This study investigated noninvasively the effects of trunk lean and foot lift exercises during sitting on abdominal
285 muscle activity. Higher TrA activity was exerted by leaning the trunk to the contralateral side and lifting the
286 contralateral foot. Furthermore, TrA contribution rate in the contralateral foot lift was similar to that in maximum
287 abdominal hollowing. As elderly people and patients with low back pain who have difficulty in consciously
288 contracting abdominal muscles can easily perform trunk lean and foot lift during sitting, these results may be useful
289 for rehabilitation to stabilize the trunks in elderly people and patients with low back pain.

290 **Compliance with Ethical Standards**

291 **Disclosure of potential conflicts of interest**

292 Conflict of Interest: The authors declare that they have no conflict of interest.

293

294 **Research involving Human Participants and/or Animals**

295 Ethics approval: All the procedures performed in the studies involving human participants were in accordance with

296 the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration

297 and its later amendments or comparable ethical standards. This study was approved by the ethics committee of

298 Kyoto University Graduate School and the Faculty of Medicine (R0546-2)

299

300 **Informed consent**

301 Informed consent was obtained from all individual participants involved in the study.

302

303 **References**

- 304 Allison GT, Morris SL, Lay B (2008) Feedforward Responses of Transversus Abdominis Are Directionally
305 Specific and Act Asymmetrically: Implications for Core Stability Theories. *J Orthop Sport Phys Ther*
306 38:228–237. doi: 10.2519/jospt.2008.2703
- 307 Aspden RM (1988) A new mathematical model of the spine and its relationship to spinal loading in the
308 workplace. 319–323.
- 309 Astfalck RG, O’Sullivan PB, Straker LM, et al (2010) Sitting postures and trunk muscle activity in adolescents
310 with and without nonspecific chronic low back pain: an analysis based on subclassification. *Spine (Phila*
311 *Pa 1976)* 35:1387–1395. doi: 10.1097/BRS.0b013e3181bd3ea6
- 312 Beith ID, Synnott RE, Newman SA (2001) Abdominal muscle activity during the abdominal hollowing
313 manoeuvre in the four point kneeling and prone positions. *Man Ther* 6:82–87. doi:
314 10.1054/math.2000.0376
- 315 Bergmark A (1989) Stability of the lumbar spine: A study in mechanical engineering. *Acta Orthop* 60:1–54. doi:
316 10.3109/17453678909154177
- 317 Bhadauria EA, Gurudut P (2017) Comparative effectiveness of lumbar stabilization, dynamic strengthening, and
318 Pilates on chronic low back pain: randomized clinical trial. *J Exerc Rehabil* 13:477–485. doi:
319 10.12965/jer.1734972.486
- 320 Bouillard K, Nordez A, Hug F (2011) Estimation of individual muscle force using elastography. *PLoS One*. doi:
321 10.1371/journal.pone.0029261


- 322 Busscher I, Van Dieën JH, Kingma I, et al (2009) Biomechanical characteristics of different regions of the
323 human spine: An in vitro study on multilevel spinal segments. *Spine (Phila Pa 1976)* 34:2858–2864. doi:
324 10.1097/BRS.0b013e3181b4c75d
- 325 Chou R, Snow V, Casey D, et al (2007) Clinical Guidelines Diagnosis and Treatment of Low Back Pain : A Joint
326 Clinical Practice Guideline from the American College of Physicians and the American. *Ann Intern Med*
327 147:478–491.
- 328 Claus AP, Hides JA, Moseley GL, Hodges PW (2018) Different ways to balance the spine in sitting: Muscle
329 activity in specific postures differs between individuals with and without a history of back pain in sitting.
330 *Clin Biomech* 52:25–32. doi: 10.1016/j.clinbiomech.2018.01.003
- 331 Ekstrom RA, Osborn RW, Hauer PL (2008) Surface Electromyographic Analysis of the Low Back Muscles
332 During Rehabilitation Exercises. *J Orthop Sport Phys Ther* 38:736–745. doi: 10.2519/jospt.2008.2865
- 333 Eriksson Crommert M, Tucker K, Holford C, et al (2017) Directional preference of activation of abdominal and
334 paraspinal muscles during position-control tasks in sitting. *J Electromyogr Kinesiol* 35:9–16. doi:
335 10.1016/j.jelekin.2017.05.002
- 336 Hasegawa K, Kitahara K, Hara T, et al (2008) Evaluation of lumbar segmental instability in degenerative
337 diseases by using a new intraoperative measurement system. *J Neurosurg Spine* 8:255–262. doi:
338 10.3171/spi/2008/8/3/255
- 339 Hides J, Stanton W, Freke M, et al (2008) MRI study of the size, symmetry and function of the trunk muscles
340 among elite cricketers with and without low back pain. *Br J Sports Med* 42:509–513. doi:

- 341 10.1136/bjism.2007.044024
- 342 Hides J, Wilson S, Stanton W, et al (2006) An MRI investigation into the function of the transversus abdominis
343 muscle during “drawing-in” of the abdominal wall. *Spine (Phila Pa 1976)* 31:175–178. doi:
344 10.1097/01.brs.0000202740.86338.df
- 345 Hill JC, Dunn KM, Lewis M, et al (2008) A primary care back pain screening tool: Identifying patient subgroups
346 for initial treatment. *Arthritis Care Res* 59:632–641. doi: 10.1002/art.23563
- 347 Hodges PW, Eriksson AEM, Shirley D, Gandevia SC (2005) Intra-abdominal pressure increases stiffness of the
348 lumbar spine. 38:1873–1880. doi: 10.1016/j.jbiomech.2004.08.016
- 349 Hodges PW, Gandevia SC, Richardson CA (1997) Contractions of specific abdominal muscles in postural tasks
350 are affected by respiratory maneuvers. *J Appl Physiol* 83:753–760.
- 351 Hodges PW, Pengel LHM, Herbert RD, Gandevia SC (2003) Measurement of muscle contraction with
352 ultrasound imaging. *Muscle and Nerve* 27:682–692. doi: 10.1002/mus.10375
- 353 Hodges PW, Richardson CA (1996) Inefficient muscular stabilization of the lumbar spine associated with low
354 back pain. A motor control evaluation of transversus abdominis. *Spine (Phila. Pa. 1976)*. 21:2640–50.
- 355 Hodges PW, Richardson CA (1997) Contraction of the abdominal muscles associated with movement of the
356 lower limb. *Phys Ther* 77:132–142; discussion 142-144.
- 357 Hodges PW, Richardson CA (1998) Delayed postural contraction of transversus abdominis in low back pain
358 associated with movement of the lower limb. *J Spinal Disord* 11:46–56. doi: 9493770
- 359 Joukar A, Shah A, Kiapour A, et al (2018) Sex Specific Sacroiliac Joint Biomechanics During Standing Upright:

- 360 A Finite Element Study. *Spine (Phila Pa 1976)* 43:E1053–E1060. doi: 10.1097/BRS.0000000000002623
- 361 Kiesel KB, Underwood FB, Mattacola CG, et al (2007) A Comparison of Select Trunk Muscle Thickness
362 Change Between Subjects With Low Back Pain Classified in the Treatment-Based Classification System
363 and Asymptomatic Controls. *J Orthop Sport Phys Ther* 37:596–607. doi: 10.2519/jospt.2007.2574
- 364 Koh HW, Cho SH, Kim CY (2014) Comparison of the Effects of Hollowing and Bracing Exercises on Cross-
365 sectional Areas of Abdominal Muscles in Middle-aged Women. *J Phys Ther Sci* 26:295–299. doi:
366 10.1589/jpts.26.295
- 367 MacDonald D, Wan A, McPhee M, et al (2016) Reliability of Abdominal Muscle Stiffness Measured Using
368 Elastography during Trunk Rehabilitation Exercises. *Ultrasound Med Biol* 42:1018–1025. doi:
369 10.1016/j.ultrasmedbio.2015.12.002
- 370 Macedo LG, Maher CG, Hancock MJ, et al (2014) Predicting Response to Motor Control Exercises and Graded
371 Activity for Patients With Low Back Pain: Preplanned Secondary Analysis of a Randomized Controlled
372 Trial. *Phys Ther* 94:1543–1554. doi: 10.2522/ptj.20140014
- 373 Masani K, Sin VW, Vette AH, et al (2009) Postural reactions of the trunk muscles to multi-directional
374 perturbations in sitting. *Clin Biomech* 24:176–182. doi: 10.1016/j.clinbiomech.2008.12.001
- 375 O’Sullivan PB, Grahamslaw KM, Kendell M, et al (2002) The Effect of Different Standing and Sitting Postures
376 on Trunk Muscle Activity in a Pain-Free Population. *Spine (Phila Pa 1976)* 27:1238–1244. doi:
377 10.1097/00007632-200206010-00019
- 378 Okubo Y, Kaneoka K, Imai A, et al (2010) Electromyographic analysis of transversus abdominis and lumbar

- 379 multifidus using wire electrodes during lumbar stabilization exercises. *J Orthop Sport Phys Ther* 40:743–
- 380 750. doi: 10.2519/jospt.2010.3192
- 381 Okubo Y, Kaneoka K, Shiina I, et al (2013) Abdominal Muscle Activity During a Standing Long Jump. *J Orthop*
- 382 *Sport Phys Ther* 43:577–582. doi: 10.2519/jospt.2013.4420
- 383 Panjabi M, Abumi K, Duranceau J, Oxland T (1989) Spinal stability and intersegmental muscle forces. A
- 384 biomechanical model. *Spine (Phila. Pa. 1976)*. 14:194–200.
- 385 Panjabi MM (1992) The stabilizing system of the spine. Part II. Neutral zone and instability hypothesis. *J. Spinal*
- 386 *Disord.* 5:390–396; discussion 397.
- 387 Richardson CA, Hides JA, Wilson S, et al (2004) Lumbo-pelvic joint protection against antigravity forces: motor
- 388 control and segmental stiffness assessed with magnetic resonance imaging. *J Gravit Physiol* 11:119–122.
- 389 Saragiotto BT, Maher ÅCG, Tie ã (2016) Motor Control Exercise for Nonspecific Low Back Pain. *Spine (Phila*
- 390 *Pa 1976)* 41:1284–1295. doi: 10.1002/14651858.CD012004.
- 391 Shiju Majeed A, Anish TS, Sugunan A, Arun MS (2019) The effectiveness of a simplified core stabilization
- 392 program (TRICCS—Trivandrum Community-based Core Stabilisation) for community-based intervention
- 393 in chronic non-specific low back pain. *J Orthop Surg Res* 14:86. doi: 10.1186/s13018-019-1131-z
- 394 Shimizu I, Tateuchi H, Motomura Y, et al (2019) Abdominal girth as an index of muscle tension during
- 395 abdominal hollowing: Selecting the optimal training intensity for the transversus abdominis muscle. *J*
- 396 *Biomech.* doi: 10.1016/j.jbiomech.2019.04.018
- 397 Tateuchi H, Akiyama H, Goto K, et al (2018) Sagittal alignment and mobility of the thoracolumbar spine are

- 398 associated with radiographic progression of secondary hip osteoarthritis. *Osteoarthr Cartil* 26:397–404.
- 399 doi: 10.1016/j.joca.2017.12.005
- 400 Tesh KM, Dunn JS, Evans JH (1987) The abdominal muscles and vertebral stability. *Spine (Phila Pa 1976)*
- 401 12:501–508.
- 402 Tsao H, Hodges PW (2007) Immediate changes in feedforward postural adjustments following voluntary motor
- 403 training. *Exp Brain Res* 181:537–546. doi: 10.1007/s00221-007-0950-z
- 404 Whittaker JL, McLean L, Hodder J, et al (2013) Association Between Changes in Electromyographic Signal
- 405 Amplitude and Abdominal Muscle Thickness in Individuals With and Without Lumbopelvic Pain. *J*
- 406 *Orthop Sport Phys Ther* 43:466–477. doi: 10.2519/jospt.2013.4440
- 407 Yamamoto I, Panjabi MM, Crisco T, Oxland T (1989) Three-dimensional movements of the whole lumbar spine
- 408 and lumbosacral joint. *Spine (Phila Pa 1976)* 14:1256–60. doi: 10.1097/00007632-198911000-00020
- 409
- 410
- 411
- 412



413


414 **Fig. 1** Task postures. a reference posture; b posterior trunk lean; c ipsilateral trunk lean; d contralateral trunk lean;

415 e ipsilateral foot lift; f contralateral foot lift; g abdominal hollowing with maximal effort. The reference posture

416 was defined as a natural posture for each participant where the perpendicular line from ear hole to the floor was

417 between the anterior and posterior superior iliac spine on the sagittal plane

418



419

420 **Fig. 2** Representative images and measurement sites of the stiffness of abdominal muscles

421

422

423

424 **Table 1** The stiffnesses of abdominal muscles and the contribution rate of transversus abdominis during tasks

	Transversus abdominis [kPa]	Internal oblique [kPa]	External oblique [kPa]	Rectus abdominis [kPa]	Contribution rate of transversus abdominis [%]
Abdominal hollowing with maximal effort (1)	39.5 ± 18.0	48.2 ± 20.9	12.4 ± 9.3	18.2 ± 8.9	33.8 ± 7.5
Reference posture (2)	11.3 ± 4.7	13.8 ± 7.9	6.8 ± 4.3	7.9 ± 3.2	28.7 ± 8.4
Posterior trunk lean at 15° (3)	12.3 ± 10.7	11.3 ± 10.1	36.3 ± 14.1	36.1 ± 19.0	12.0 ± 7.7
Posterior trunk lean at max (4)	9.3 ± 4.3	10.7 ± 4.8	66.4 ± 21.1	70.6 ± 22.0	6.1 ± 2.8
Ipsilateral trunk lean at 20° (5)	13.3 ± 6.4	14.3 ± 6.0	8.0 ± 4.9	18.0 ± 17.7	25.9 ± 8.9
Ipsilateral trunk lean at max (6)	18.8 ± 9.4	20.5 ± 10.1	12.8 ± 6.9	20.1 ± 10.2	25.8 ± 7.0
Contralateral trunk lean at 20° (7)	19.6 ± 8.1	26.5 ± 10.6	42.1 ± 10.3	13.7 ± 6.2	19.5 ± 7.7
Contralateral trunk lean at max (8)	26.1 ± 11.5	36.1 ± 15.3	55.4 ± 19.0	27.7 ± 18.4	18.3 ± 7.0
Ipsilateral foot lift (9)	14.0 ± 4.6	15.4 ± 6.4	9.6 ± 7.0	9.2 ± 5.7	29.3 ± 7.7
Contralateral foot lift (10)	18.0 ± 7.8	20.7 ± 7.7	7.1 ± 7.1	13.6 ± 11.4	31.2 ± 8.8

425 Values are expressed as mean ± standard deviation

426 ¹⁻¹⁰ *P* < 0.05 vs. the task, which is corresponded to numbers

427 **Table 2** Spinal alignment during each task

	Thoracic kyphosis [°] (n=12)	Lumbar lordosis [°] (n=12)	Pelvic inclination [°] (n=14)	Spinal inclination [°] (n=14)
Reference posture	29.1 ± 6.0	2.8 ± 7.3	1.0 ± 9.1	
Posterior trunk lean at 15°	31.0 ± 6.1	4.0 ± 9.6	11.9 ± 8.8 *	
Posterior trunk lean at max	32.0 ± 6.4	5.8 ± 8.0	24.8 ± 11.5 *	28.7 ± 5.2
Ipsilateral trunk lean at 20°	34.0 ± 8.2	10.9 ± 10.8	-1.5 ± 9.6	
Ipsilateral trunk lean at max	29.6 ± 6.8	8.2 ± 6.4	-1.8 ± 8.9	28.5 ± 3.8
Contralateral trunk lean at 20°	31.2 ± 6.4	8.7 ± 7.6	-1.4 ± 9.5	
Contralateral trunk lean at max	31.1 ± 6.8	6.8 ± 6.9	-4.2 ± 8.7	29.4 ± 4.6
Ipsilateral foot lift	28.2 ± 5.9	7.9 ± 8.4	1.7 ± 10.1	
Contralateral foot lift	26.8 ± 6.0	7.1 ± 6.5	0.2 ± 9.9	

428 Values are expressed as mean ± standard deviation

429 The positive values in pelvic inclination represent the sacral posterior inclination angle on the sagittal plane

430 * $P < 0.05$ vs. reference posture

431